

PEEL CORNER
OF BOOKLET

5M-7-29

Released For Shipment By:
VOLUNTARY PURCHASING GROUPS, INC.
230 FM 87
BONHAM, TEXAS 75148
EPA Reg. No. 7401-311 • EPA Est. No. 7401-TX-1
Visit Us At: www.fertilome.com



ferti·lome[®]



Fire Blight Spray

- For Apples, Pears, Pyracantha, Chrysanthemums, Philodendron, Dieffenbachia, And Roses.
- 1 Tablespoon Makes 2 1/2 Gallons Of Spray.

ACTIVE INGREDIENTS:

Streptomycin Sulfate . . .21.2%;

INERT INGREDIENTS: . . .78.8%

*Contains 17% Streptomycin derived from 21.2% Streptomycin Sulfate.

Keep Out Of Reach Of Children

CAUTION

See Back Panel For Additional
Precautionary Statements

NET CONTENTS 2 Ozs.

3. HERE ARE THE RESULTS YOU SHOULD EXPECT!

When used according to directions, this product will control *Fire Blight* and *Bacterial Wilt, Stem Rot, Leaf Spot, and Crown Gall* on Fruit Trees, Flowers and Shrubs.

STORAGE AND DISPOSAL

STORAGE: Store in original container and place in a locked storage area.

DISPOSAL: Do not reuse container. Wrap and put in trash.

Released For Shipment By:
VOLUNTARY PURCHASING GROUPS, INC.
230 FM 87 • BONHAM, TEXAS 75418
EPA Reg. No. 7401-311 • EPA Est. No. 7401-TX-1
Visit Us At: www.fertilome.com



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

May cause allergic skin reactions. Do not breathe dust or spray mist. Wear a dust mask and rubber gloves. Wash thoroughly after handling. This material is not to be used for medical, veterinary, or human purposes. DO NOT APPLY this product in a way that will contact unprotected workers, either directly or through drift. Only protected handlers may be in the area during application. DO NOT ENTER or allow entry into treated areas until (sprays have dried/dusts have settled/vapors have dispersed, as applicable) to perform hand labor tasks.

STATEMENT OF PRACTICAL TREATMENT

If the pesticide comes in contact with skin, wash off with soap and water. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or toileting. Before removing gloves, wash them with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste. *Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

1. WHAT IS THIS PACKAGE FOR?

Use this product to control *Fire Blight and Bacterial Wilt, Stem Rot, Leaf Spot, and Crown Gall* on Fruit Trees, Flowers and Shrubs.

2. HOW DO YOU USE IT?

FRUIT: APPLES AND PEARS (Areas Other Than West Coast): To control *Fire Blight* in Home Plantings, apply 1 tablespoon in 2½ gallons of water (equivalent to 100 ppm or 4 ozs. per 50 gallons of water) in foliar and blossom sprays. Make first spray at the start of blossoming period. Continue spray applications every 3 to 4 days during bloom time. Apply additional sprays every 5 to 7 days after blossom period when weather conditions favor spread of Fire Blight. Do not apply when fruit is visible.

PYRACANTHA: To control *Fire Blight*, apply a solution of 1 level tablespoon of **ferti-lome® FIRE BLIGHT SPRAY** per 2½ gallons of water (100 ppm concentration). Begin spraying at start of blossom period. Repeat at 3 to 4 day intervals during bloom and at 5 to 7 days intervals after bloom if weather favors Disease spread.

CHRYSANTHEMUMS (Cuttings): To control *Bacterial Wilt*, soak cuttings for 4 hours in a solution containing ½ tablespoon of

ferti-lome® FIRE BLIGHT SPRAY per 2½ gallons of water (50 ppm concentration) and plant as usual.

DIEFFENBACHIA: To control *Bacterial Stem Rot*, spray plants at 5 to 7 day intervals with a solution of 1 level tablespoon of **ferti-lome® FIRE BLIGHT SPRAY** in 2½ gallons water (100 ppm concentration) or soak cuttings for 20 minutes in a solution of 2 level tablespoons in 2½ gallons of water (200 ppm concentration) and plant in sterile rooting medium.

PHILODENDRON: To control *Bacterial Stem Rot*, remove all decayed leaves and spray with a solution of 2 level tablespoons of **ferti-lome® FIRE BLIGHT SPRAY** in 2½ gallons of water (200 ppm concentration) as preventative or at first sign of water soaked leaves. Repeat at 4 to 5 day intervals.

ROSES: To control *Crown Gall*, remove infested Plant and cut out gall tissue. Soak root system and cut surfaces of infested area for 15 minutes in a solution of 2 level tablespoons of **ferti-lome® FIRE BLIGHT SPRAY** per 2½ gallons water (200 ppm concentration). Replant in soil free from Disease organisms, and apply ½ tablespoon per 2½ gallons water (50 ppm concentration) as foliage spray at weekly intervals.