

**CONCENTRATE**  
Makes 4 Gallons

# Garden Fungicide

**FOR FLOWERS, FRUIT & VEGETABLES**

**Safer**  
BRAND

• Controls  
Powdery  
Mildew,  
Black Spot  
& Rust

ACTIVE INGREDIENT  
Sulfur..... 12.00%  
OTHER INGREDIENTS..... 88.00%  
TOTAL..... 100.00%

KEEP OUT OF REACH OF CHILDREN

**CAUTION** See back panel for additional  
Precautionary Statements  
and First Aid.

www.victorpest.com

NET CONTENTS 16 FL. OZ. (473 ml)



**CONCENTRATE**

# Garden Fungicide

**FOR FLOWERS, FRUIT & VEGETABLES**

**Safer**

**Always Shake Well Before Using**

### DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Shake well. For control of **Blackspot, Leaf Spot, and Rust**, mix 1 part concentrate to 30 parts water or 4 fl. oz. (8 tbsp.) per gal. of water. For **Powdery Mildew** mix 2 fl. oz. (4 tbsp.) per gallon of water. Place about half of the volume of water in sprayer; add concentrate; stir well; add rest of water and mix thoroughly. Run agitator on large tank sprayers. For coverage see individual applications. Spray plants thoroughly to wet the upper and lower surfaces of foliage. Repeat as recommended. To control powdery mildew, leaf spot, and rust on **Flowers, Foliage Plants & Ornamentals**, spray at 7- to 10-day intervals or as needed. To control powdery mildew, black spot and rust on **Roses**, spray at 7- to 10-day intervals, or following rain. Be sure to treat new growth & buds. Avoid spraying when plants are in full sun or temperature exceeds 85°F. Apply 1 gal. mix to each 350 sq. ft. To control powdery mildew on **Beans & Peas**, spray at first sign of disease & repeat at 7-day intervals as necessary. May be used up to day before harvest. Apply 1 gal. of mix to each 250 sq. ft. To control powdery mildew on **Apples & Pears**, apply to thoroughly wet foliage. Apply at pre-bloom stage or after petal fall as needed. **Note:** Some apple & pear varieties are sulfur sensitive; test for varietal sensitivity prior to full scale treatment. To control powdery mildew on **Potatoes**, spray before disease appears & repeat as necessary. Apply 1 gal. of mix to each 350 sq.

ft. On **Cucumbers and Squash**, spray when powdery mildew appears & repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 600 sq. ft. To control powdery mildew on **Strawberries**, spray when disease appears and repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 300 sq. ft. **Note:** Do not apply to strawberries that will be used for canning. May be used up to the day before harvest. To control powdery mildew on **Grapes**, thoroughly cover foliage. Begin when new shoots are 6-10 inches. Repeat before blooms open and at 14-day intervals as needed. **CAUTION:** Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord & other bunchless varieties. **Home Greenhouses:** To control powdery mildew, leaf spot and rust on bedding plants, propagates, fruits & vegetables: Mix 1 part concentrate to 30 parts water (8 tbsp. per gal.) and apply according to above recommendations by crop type. Test for phytotoxicity before full-scale application. **Use Cautions:** Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not apply on Boston fern, spinach, arcticans, filberts, walnuts, and yucca. Do not apply when temperature exceeds 85°F. Do not apply for at least 4 weeks following application of an oil spray or within 4 days of other pesticides. Be sure to wash and rinse spray equipment thoroughly after each use. Do not allow mixed spray to stand in spray equipment. **Re-entry:** Avoid contact with plants until spray has dried.

**ENVIRONMENTAL HAZARDS:**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of rinsale or equipment washwater.

### STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal. **Storage:** Store in a cool place. **Dispose:** If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**Woodstream Corp. and its subsidiaries will not accept liability for damage or injury resulting from misuse. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.**

Distributed By: Safer<sup>®</sup>, Inc., a wholly-owned subsidiary of Woodstream Corporation

**Woodstream**

89 North Locust Street, Litz, PA 17543

1-800-900-1819 • 717-626-2125

EPA Reg. No. 42697-37

EPA Est. No.

Ⓜ 44616-MO-001 Ⓜ 11600-MN-001

Ⓜ 63416-MN-001

Circled letter corresponds to first letter of lot number on label, container side or bottom.

Look for other Safer<sup>®</sup> Brand products at [www.victorpest.com](http://www.victorpest.com)

© Copyright 2002 Woodstream Corp.

5456-1549-RMS456B Model No. 5456

Made in USA

