

BIONEEM®

INSECTICIDE & REPELLENT



MULTI-PURPOSE CONCENTRATE

NET CONTENTS 16 FL. OZ. (473 ml)

For Fruit, Vegetables,
 Flowers & Ornamentals
 In & Around the Home

◆ Kills or Repels Aphids, Beetles, Caterpillars, Leafminers, Thrips, Whiteflies and other listed pests.

KEEP OUT OF REACH OF CHILDREN

CAUTION See back panel for additional Precautionary Statements and First Aid.

0204

GENERAL INFORMATION:

BIONEEM® effectiveness is based on several modes of action. It disrupts insects' delicate hormonal balance so they de before they most to the next life stage. BIONEEM® suppresses some insects' desire to feed adso repels many insects in areas that have been sprayed, hence different modes of action result in braid spectrum control of the majority of insects that harm your yard and garden. BIONEEM® risk for repels a broad ascortment of insect pests. BIONEEM® sprace many beneficial insects what so noney bees and lady bugs that prey on undesirable insects. Without beneficial insects to control pest populations, plants are even more vulnerable to new intestations.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Stake well before using. Use spray solutions within 8 hours of mixing for maximum effectiveness. Do not store diluted spay solution for later use. Not for use in food handing mass. Since BOINEEM® interferes with rissect growth function, it is most effective when applied 2-3 times at 7-10 day intervals. Application directly onto the target pest will increase efficacy. Time applications for early to mixi-moming or in the late afternoon. Do not add other chemicals to BIONEEM® solutions. Do not apply to writed, stressed, or nen'ny transplanted materials prior to not development. Do not apply to known sensitive plant species without prior testing.

Known Sensitive Plants:

Do not apply BIONEEM® to the following plant species due to potential plant damage: Hibiscus flowers

Foliar Application (Bedding & Potted Plants): For use on bedding plants, sorted and foliage plants, ornamentals, trees and shrusts, in and around forests to control the immuture or leval forms of the following insect pests: Leaflingpers, Leafminers, Whitefiles, Thrips, Mealybugs, Loopers, Caterpillars, Aphids, Fruit Illes, Beet Armyworms, Mexican Bean Beetles. USE DIRECTIONS: Dilute SIDNERS¹¹ to a concentration of 3 th. Cz. (6 balesponos) of BIONEEM¹⁹ cagnion of vater (1 part BIONEEM¹⁹ to 40 parts vater), Mix thoroughly, Apply with a hand sprayer as a fine spray to leaf surfaces to run-off. Be sure to coat undersides of leaves as this is where many insects bids. And decessive accollates.

Foliar Application (Trees & Shrubs): For use on trees and shrubs on residential landscapes to control the immuture laval stages of the following insect pests: Leafhoppers, Leafminers, Oppsy moths, Weevils, Psyllids, Webworms, Hornworms, Spruce budworms, Pine sawflies, Elm leaf beetle, Bagworms, Tent Caterpillars, Grasshoppers. USE DIRECTIONS: Dilute BIONEEMP to a concentration of 3 ft. 0z. (6 tablespoons) of BIONEEMP or application of water (1 part BIONEEMP to a polymer water). Mix throughly, Apply with a hand sprayer as a fine spray to leaf surfaces to run-off. Be sure to coat undersides of leaves as this k where many insects hide, Auch docssesse anolitical coater.

Lawn Application: For home lawns to control the following insect pests: Mole crickets, White grubs, Sod webworms. Fall armyworms. Chinch bugs. Hyperodes weevils. Billbugs. Crane files.

USE DIRECTIONS: Dilute 2 ft. Oz. (4 tablespoons) of BIONEEM® in 1 gallon of water. Apply at the rate of 5 gallons spray solution per 2000 square feet. Apply BIONEEM® 2-3 times on a 7-10 st interval to insure effective control of surface and sub-race feeding pests. Do not apply BIONEEM® to newly seeded lawns. Best timing for initial white grub application will vary by region but is generally early to mid summer and again in the fall. It in doubt, contact your local university research or extension specialist for the periods when which orous are feeding in your area.

Yard and Landscape BIONEEM® Concentrater. For use around the home to control the following insects: Aphids, Caterpillars, Gypsy moth larvae, immalure tent beetles, Webworms, Bagworms, Lace bugs, Box elder bugs, Leathoppers, Psyllids, Sawtly larvae, Wooly aphids. USE DIRECTIONS: Always use freshly mixed spray solution. Use soft water whenever possible. Use with caution on new tender foliage, and do not apply to plants in direct sunlight. When insects are first noticed, apply a mixture of 3 ft. Dz. (6 tablespoons) of BIONEEM® per gallon of water (1 part BIONEEM® to 40 parts water). Repeat applications 2-3 times on a 7-day interval. When intestations are heavy, apply on a 3-4 day interval until populations are refused.

House Plant Application: For use on house plants to kill or repel the following insect pests: Leafhogners, Leaf miners, Whitellies, Thrips, Mealybugs, Loopers, Caterpillars, Aphids, Fruit files, Beet armyworms, Mexican bean beetles, Fruguss gnats. USE DIRECTIONS: Dilute BIOHESM[®] to a concentration of 0.4.2 to 2, 12 teaspoons) of BIONESM[®] per pint of water. Mix thoroughly, Aglust spray bottle nozice to spray setting and apply BIONEEM[®] be lost surfaces until completely wet Ee surf to cast understand. leaves especially those at the soil surface, as this is where many insects hide. Avoid excessive application.

Vegetable and Fruit Crop Application: For use on vegetable and fruit crops to kill or repel the following insect pests: Armyworms, Diamondback Moths, Hornworms, Imported cabbageworms, Loopers, Melonworms, Pickleworms, Tomato fruitworms, Corn earworms, Bollworms, Budworms, Whiteflies, Thrips, Borers, Aphids, Squash bugs, Tent caterpillars, Mexican bean beetles, Pinworms, Colorado notato heetles, Furonean com horers, Cucumber heetles, Leaf miners, Vine horers, Cutworms. Weevils, Bean leaf beetles, Leafrollers, Grape leaf skeletonizers, Twig girdlers, Codling moths on the following crops: Broccoli, Turnips, Cabbage, Tomatoes, Melons, Kale, Mustard, Peppers, Lentils, Squash, Lettuce, Beans, Corn. Potatoes, Peanuts, Asparagus, Garlic, Parsley, Radishes, Rutabaga, Anise, Mint, Dill, Rosemary, Sweet bay, Thyme, Almonds, Pecans, Pistachios, Apples, Pumpkins, Peaches, Plums, Figs, Lemons, Cauliflower, Celery, Kohlrabi, Carrots, Spinach, Turnip greens, Brussels sprouts, Sweet potatoes, Cucumbers, Collards, Beets, Peas, Egoplants, Ginger, Artichokes, Onions, Turnips, Bok chov, Collard greens, Mustard greens, Basil, Chives, Marioram, Sage, Tarragon, Wintergreen, Cashews, Macadamia nuts, Walnuts, Cherries, Cantaloupes, Pears, Apricots, Oranges, Limes, USE DIRECTIONS: Always use freshly mixed spray solution. Use soft water whenever possible. Use with caution on new tender foliage. When insects are first noticed, apply a mixture of 3 fl. Oz. (6 tablespoons) of BIONEEM® per gallon of water (1 part BIONEEM® to 40 parts water). Repeat applications 2-3 times on a 7-day interval. When infestations are heavy, apply on a 3-4 day interval until populations are reduced

Repellency Against Japanese Beetles and Other Pests:

BIONEEM® regists a variety of pests including armynorms, aphids, and Japanese Beetles. Dilute BIONEEM® to a concentration of 3 ft. 02. (6 biblespoons) of BIONEEM® per gallon of water (1 part BIONEEM® to 40 parts water). Mix thoroughly. Apply to plants with a hand sprayer, being sure to completely cover foliage and flowers. Apply just before pests emerge or at first sign of infestation. Repeat apolication server 3-2 days or immediately after significant rainfall.

PRECAUTIONARY STATEMENTS:

Hazards to Humans and Domestic Animals:

Caution: Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Harmful if inhaled. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything to an unconscious person, **If on Skin or Clothing**: Take off contaminated clothing, Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If in Eves: Hold eve open and rinse slowly and gently with water for

15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible, Call a poison control center or doctor for treatment advice Have the product container or label with you when calling a poison control center or

doctor, or going for treatment. Hot Line Number: 1-800-858-7378.

ENVIRONMENTAL HAZARDS:

This product is hazardous to fish and aquatic invertebrates. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS:

Flammable. Keep away from heat and open flame.

STORAGE AND DISPOSAL:

Pesticide Storage: Keep in original container. Store in a cool, dry place, away from direct sunlight, feed or foodstuffs. Keep container tightly sealed when not in use. Do not store below 50°F (10°C) or above 95°F (35°C).

Pesticide and Container Disposal:

If Empty: Do not reuse this container, Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Woodstream Corp. and its subsidiaries will not accept liability for damage or injury resulting from misuse. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Distributed By: Safer®, Inc., a wholly-owned subsidiary of Woodstream Corporation

₩woodstream

69 North Locust Street, Lititz, PA 17543 800-800-1819 • 717-626-2125

EPA Reg. No. 70051-6-42697 EPA Est. No. 63456-MN-001

Look for other Safer® Brand products at www.saferhrand.com



© Copyright 2005 Woodstream Corp. 5612-3320-RM06395 Model No. 5612 Made in USA

PF 6926

0305

BIONEEM® INSECTICIDE & REPELLENT Safer

ShakeWell Before Using!

MULTI-PURPOSE CONCENTRATE

Woodstream Corp. and its subsidiaries will not accept liability for damage or injury resulting from misuse. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-838-7378.

Distributed By: Safer®, Inc., a wholly-owned subsidiary of Woodstream Corporation

₩woodstream

69 North Locust Street, Lititz, PA 17543 800-800-1819 • 717-626-2125 EPA Reg. No. 70051-6-42697

EPA Est. No. 63456-MN-001

Look for other Safer® Brand products at www.saferbrand.com

0305



PULL HERE TO OPEN / PRESS TO RESEAL ▶

© Copyright 2005 Woodstream Corp. 5612-3320-RM06395 Model No. 5612 Made in USA