


RED LINE INDICATES DIE - DOTTED RED LINE SHOWS LEAFLET PLACEMENT

BLUE LINES SHOW WHERE COPY AND GRAPHICS CAN BE PLACED - SHADED BLUE AREA WILL BE UNDERNEATH THE LEAFLET

YELLOW AREA INDICATES HOW FAR COLOR/GRAPHICS SHOULD BLEED OUTSIDE DIE



RED LINE INDICATES DIE - DOTTED RED LINE SHOWS LEAFLET PLACEMENT

BLUE LINES SHOW WHERE COPY AND GRAPHICS CAN BE PLACED - SHADED BLUE AREA WILL BE UNDERNEATH THE LEAFLET

YELLOW AREA INDICATES HOW FAR COLOR/GRAPHICS SHOULD BLEED OUTSIDE DIE

COPY AREA PAGE 1

COPY AREA PAGE 2

COPY AREA PAGE 3

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product can be applied with all types of hand or power sprayers. Read and carefully follow the instructions for the type of sprayer you use.

used up to and including Management (IPM) progra SMAKE WILL, for best results, use frestly mixed solution Use soft water witermer prossible. NOTE: Do not use on new transplants, newly coded cuttings or plants stossed by frought. World application when temperature exceeds 90°F. Stafe if Brand Insect Killing Stop may be used up to and incultuing day of harvest, and is useful as a component of integrated Pest Management (PM) programs.

Indoor Plants: Mix 2.5 ft. oz. (§ Tlosp.) of concentrate per gallon of water. Apply when freects or signs of their damage appear. Take care to thoroughly wet all surfaces of interest of longs and branches for Anthats, seamine paints after two days and spary again in freeded. For other pests, sprary wesley at 7 to 10-day intervals as needed up to three times. Do not make repeated snrays a stoot intervals without throughly instead polary, for 0 testors. There is no pests sprary wesley at 7 to contact things on the blooms. Root Mealyhous and Springfall repeated snrays as door iterative on the two to belspoons of concentrate in a fine station of the or electron station of the original minestations can be reduced using a soll depend on the two to belspoons of concentrate in a quart of the original per soll resolution. Do not use on faceing and stationary supports the contraction of the original per soll resolution of the original per soll resolution. On one use on faceing deficient electric differentiation, originate that was, paints or succulents without testing.

Toward and Bedding Plants Coroning Outdoors and in Home Greenhouses: Including foliage points: roses and subculents. May 25 ft. oz. (5 18p.) of concentrate per gation of water. Apply dentis: roses and subculents. May 25 ft. oz. (5 18p.) of concentrate per gation of water. Apply when the person of the perso bleofting heart (gardenias, jade plant, lantana, Illies or sweet peas. Some varieties of azaleas begonias, camellias, fuschias, gardenias and impatiens have shown sensitivity. If withing locurs within a few hours, rinse with clean water supply. Test on chrysanthemums for varieta

Unannernal Trees and Shruber, Max 2.5 ft. oz. (5 10sp.) of concentrate per gallon of wetter. To control the crawler stage of Medigids, Psyllidis and Seales, spray to we butk surface when intestation warrants. For Mittage is part to wet underside of leaves and medical seaths. For intestation warrants. For Mittage is part to wet underside of leaves and medical seaths. For intestation warrants. For Mittage is present. Do not apply to everguents during drought or whom beneficially as present. Do not apply to be express during drought or whom congregated on ordered of them. Write The not apply to observed, sealeres raugh more than congregated on ordered of them. Write the notation of the sealer of Colorado bius sproze. Do not apply to observed, sealeres raugh more than a congregated on ordered of the medical and drought establishes statistically sealer to the sealer of Colorado bius sprozes. Do not apply to observed, sealeres Raught more sprozes. The sealer of the sealer o

Legume Vegetables: Including beans, peas and soybeans to control aphids, grasshoppers whiteflies and spider mites. Use up to day of harvest.

Frailing Vegladbest. Including formations, peoples and egiplant to control spider mites and writely and to suppress apiniss. Use up to day of harvest.

Countri Vegladbest including quounitier, summer squasts and melon to control plant bugs and writely and to suppress apiniss and mites. Use up to day of harvest. Always apply to promotives a triplat or funding the contest part of the day.

Identical spices: Including basil, chives, dill, marjoram, and sage to control writelity and to spices. Including basil, chives, dill, marjoram, and sage to control writelity and to

suppress admities, bits up to day of hanvest.

Finite and Must, Mik 2.5 no. (5 Tisps) of concentrate per gallon of water. Spray when
theses first space or to present damage, Apply to achieve through coverage of infester
foliage and/or bandness. Repeat at weekly internats as needed, Avoid spraying when
besome and present. Use up to day of harvest.

Drius Fruits: Including lerroins, sueel crainipe and graphrist to control achies, cottony bushion scale, back scale, red scale and miles. Use up to Gay of francest. NOTE. Some marking of fluid may occur, especially when spay roundi occurs on when soody mold is present. Under Fruits: Including apple and peer to control leaffragores, scale crawins, paer goyla, paer stag, fleminariale) and to suppress against and marks. Use up to day of lancest. MICTE. paer stag, fleminariale and to suppress against and marks. Use up to day of lancest with red party may cause localized marking of some varieties, particularly apple varieties such as Red and Goden pelicious, Grameristin, and pear varieties south as d'hique, Compris and Asian. Stone Fruits: Including cherry, peach and Jumis to control scale insects and suppress points and miles. Les up to day of the marks. MICTE. Spray may cause localized sporting of markships and enrove waveform from plums.

meanines automotive recommendation of the commentary to small final s and Berinse including blackburry, coffee, cramberry, grape and strawberry to commol learnappers, mealyburgs, scales and whitefully and to suppress aghicts and mines, they up to day or framests. URLE: Do not soy request during the period between the start of blooming and 20 3 days post starter.

There hits: tracking amounts, English walmus and peaces to control scale crawlers and in upopress aphicls and mites. Use up to day of harvest.

COPY AREA PAGE 4

PAGE 5

FRONT COVER COPY AREA

COPY AREA

ENVIRONMENTAL HAZARDS:

Inits product may be hazardous to aqualic invertebrates. For terrestrial uses, do not apply limits product may be hazardous to aqualic invertebrates. For intendial areas below tirectly to water, or to areas where surface water is present or to intendial areas below the mean high water mark. Do not contaminate water by cleaning of equipment or

Flammable. Keep away from heat a STORAGE AND DISPOSAL: PHYSICAL OR CHEMICAL HAZARDS: neat and open flame.

Hazards is Humans and Domestic Animals:

WARNING: Causes sin irritation. Causes substantial but temporary eye injury. Harmful if warning for the present on additing. Avoid absorbed through sich Harmful if initiated to not get on skin, eyes or on additing. Avoid bearanting story mist. Wash throughly with soap and water after handling and betree sating, diricking, cleaving quin, or using tobacco. Remove and wash confirmalinated bothing before reuse. West protective eyewear (goggles, face shield or safety glasses).

PRECAUTIONARY STATEMENTS:

Do not contaminate water, feed or foodstuffs by storage and disposal.

STORAGE: Store only in original container in a cool, dry area inaccessible to children and pets. Do not store full or partial containers in direct sunlight, Keep container tightly sealed

DISPOSAL: If Empty: Do not reuse this container: Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or (800)-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If Inhaled: Move person to fresh air, if person is not breathing, call 911 or an ambulance, then give artificial respiration, pretrably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going tor treatment. You may also contact the National Pesticide Information Center at 1-500-858-7378 for emergency medical treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call

poison control center or doctor for treatment advice.

Woodstream Corp. and its subsidiaries will not accept liability for damage or injury resulting from misuse. For information on this pecticide product (including health concerns, medical energencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Sise" Plard Insect (Illing Saap Concentrals effactively controls: Adeldus, Meakhous, Kent Caterplians, Aprilds; Perd Bugs, Things, Enroyls, Shaffler, Methingh, Gerschoppers, Sawhy Lannas, Woody Apolisis, Lazebusy, Sizales, Ledinoppers, Solder Mites, Safer® Gand Insect (Illing Seap Concentrate offers effective control on: Foliage Plants, Trees & Shrubs, Tree Nuts, Fower, Fruits, Wegetables.

Woodstream Corporation Distributed By: Safer^e, Inc., a wholly-owned subsidiary of

Wwoodstream

69 North Locust Street, Littz, PA 17543 1-800-800-1819 • 717-626-2125

EPA Reg. No. 42697-60 EPA Est. No. 44616-MO-001 _ook for other Safer® Brand

products at www.saferbrand.com

© Copyright 2004 Woodstream Corp. 5118-2588-RM5118 Model No. 5118 Made in USA

JOB #0000

RED LINE INDICATES DIE YELLOW AREA INDICATES HOW FA BLUE LINES SHOW WHERE COPY AND GRAPHICS CAN BE PLACED R COLOR/GRAPHICS MAY BLEED ON FRONT COVER