

CITRUS, FRUIT & NUT ORCHARD SPRAY CONCENTRATE

Controls a wide range of fungal diseases and insects on listed fruit, nuts, citrus, vegetables, ornamentals, houseplants and lawns.

Dilute, per label instructions, as little as 2 ½ oz. per gallon of water in drier weather or when low levels of disease are present.

One pint makes up to 6.4 gallons of finished spray.

Manufactured under a license of W. Neudorff GmbH KG, Germany.



EPA Est. No. 4-NY-1
EPA Reg. No. 67702-17-4

Distributed by
Bonide Products, Inc.
6301 Sutliff Road
Oriskany, NY 13424



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Shake well before using.

FOR CONTROL OF FUNGAL DISEASES

When high levels of disease are present, or during periods of wet weather, mix 5 fluid oz product with 1 gallon of water and shake well. During drier weather or when low levels of disease are present, mix 2.5 fluid oz product with 1 gallon of water and shake well. Thoroughly spray all areas of the plant, especially new shoots. For best control apply as a protective spray early in the season before the diseases are noticed. Re-spray every 7 to 10 days or after rain. May be used up to day before harvest.

FOR CONTROL OF INSECTS/MITES

Dilute product by mixing 5 fluid oz product with 1 gallon of water. Shake well. Thoroughly spray all areas of the plant, especially new shoots and the underside of leaves. Pests need to be contacted by the spray to be controlled.

USE SITE- INDOOR, OUTDOOR AND GREENHOUSE	PEST	DISEASE
FRUIT AND NUT TREES: almond, apple, avocados , cherry, citrus (such as, oranges, grapefruit, lemon, limes, citron, kumquat), filbert, nectarine, peach, pear, pecans, pistachios, plum, prune, walnut	aphids, mites, spider mites, leafhoppers, caterpillars, whiteflies, spittlebugs, mealybugs, scale, thrips, psyllids, plant bugs, fruit flies, earwigs	scab, powdery mildew, rust, blight, brown rot
VEGETABLE AND VINE PLANTS: beans, beets, blackberries, blueberries, broccoli, brussel sprouts, cabbage, carrots, cauliflower, corn, cucurbits (cucumber, melons, pumpkins and squash), currant, gooseberries, grapes, kale, onions, peas, peppers, potatoes, raspberries, strawberries, tomatoes	aphids, mites, spider mites, leafhoppers, caterpillars, whiteflies, spittlebugs, mealybugs, scale, thrips, psyllids, plant bugs, fruit flies, earwigs	powdery mildew, rust, blight, leaf spots

USE SITE- INDOOR, OUTDOOR AND GREENHOUSE	PEST	DISEASE
ORNAMENTALS: flowering plants and shrubs, woody ornamentals, roses, trees and houseplants	aphids, mites, spider mites, leafhoppers, caterpillars, roseslugs, whiteflies, spittlebugs, mealybugs, scale, thrips, psyllids, plant bugs, lacebugs, earwigs	powdery mildew, rust, blight, blackspot, leaf spots
LAWNS	aphids, mites, spider mites, leafhoppers, caterpillars, spittlebugs, thrips, chiggers, earwigs	rust, leaf spot, blight, powdery mildew

For use only on home lawns and gardens. Not for use on agricultural establishments growing agricultural crops or commodities for resale.

Note: Wet weather promotes plant diseases.

Proper plant nutrition and removing dead or damaged leaves will also reduce disease and insect problems.

USE PRECAUTIONS

- Do not use if temperatures are expected to be over 90°F.
- Do not spray plants with this product within 21 days of an oil spray.
- Some varieties of grapes and cucurbits are sensitive to sulfur and some plant damage may occur.
- Use care or do not spray on sulfur sensitive plants. If not sure, test spray a few leaves 2 days before spraying the entire plant.
- Do not apply to strawberry fruit that will be used for canning.

STORAGE AND DISPOSAL

Store at room temperature and protect from freezing.

CONTAINER DISPOSAL:

If empty: Nonrefillable. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID

IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

ENVIRONMENTAL HAZARDS

This product may be hazardous to aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather or when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

Buyer assumes all responsibility for safety and use not in accordance with directions.