

Safer[®]
BRAND

3-in-1

GARDEN SPRAY II



**Kills
Fungus**



**Kills
Mites**



**Kills
Insects**

Active Ingredients:	By Wt.
Potassium Salts of Fatty Acids.....	0.75% [†]
Sulfur.....	0.40%
INERT INGREDIENTS.....	98.85%
Total.....	100.00%

[†] Includes 0.55% as potassium oleate and 0.20% as potassium salts of other fatty acids.

OMRI[™]
Listed
Organic Materials
Review Institute

FOR
organic
gardening

**KEEP OUT OF REACH
OF CHILDREN**

See back panel for additional
precautionary statements.

NET CONTENTS 24 FL. OZ.

1004

PF-3330

WB F w/Lam



Safer
BRAND

3-in-1 GARDEN SPRAY II

Always Shake Well Before Using!

Safer® Brand 3-In-1 Garden Spray can be used to kill insect and mite pests and to prevent and control fungal disease. Safer® Brand 3-In-1 Garden Spray contains sulfur and insect killing soap derived from natural fats and plant oils. Its formulation is gentle on most plants. Safer® Brand 3-In-1 Garden Spray is designed to kill pests and create an environment where damaging fungi cannot live. The ready-to-use formulation can be applied directly to fruits and vegetables right up to the day before harvest.

PRECAUTIONARY STATEMENTS: Environmental Hazards: This product may be hazardous to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment washwaters or disposal of rinsate.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WHAT IT IS: Safer® Brand 3-In-1 Garden Spray provides three garden solutions in one: fungicide, insecticide and miticide. It kills garden insects and mites and controls fungal diseases. It contains a patented formulation to help reduce injury to most plants.

WHAT IT CONTROLS: Powdery mildew, black spot, leaf spot, and rust.

WHAT IT KILLS: Aphids, Beetles (leaf-feeding), Caterpillars (leaf-feeding), Crickets, Earwigs, Lace Bugs, Leafhoppers, Mealybugs, Mites, Plant Bugs, Scale Crawlers, Thrips, and Whiteflies.

WHERE TO USE IT: Roses, Flowers, Shade Trees, Foliage Plants, Ornamentals, Beans, Peas, Broccoli, Cabbage, Cauliflower, Collards, Kale, Peppers, Strawberries, Apples, Pears, Citrus, Potatoes, Cucumbers, Squash, Grapes, Blackberries, Blueberries, Raspberries, Loganberries, Currants, Melons, and Tomatoes.

HOW TO USE IT: Apply in the morning or late evening. Shake Well. Use at first sighting of insects or infestation. Thorough coverage of both top and bottom leaves is important. Spray leaves to the point of runoff. Apply every 5 to 7 days when insects are present.

USE CAUTIONS: Some plants are susceptible to injury from sulfur under certain climatic conditions. DO NOT USE on delicate fern, apricots, filberts, stock, walnuts, nasturtium, sweet pea, violet, mountain ash, Japanese maple, horse chestnut, spinach, viburnum, newly planted cuttings, or plants stressed by drought. Do not apply when plants are in full sun or when temperature exceeds 90° F. Do not apply for at least 4 weeks following application of oil spray.

FUNGICIDE USE: FLOWERS AND ORNAMENTALS: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain or as needed to control powdery mildew, leaf spot, black spot, and rust. Be sure to treat new growth and buds.

BEANS, PEAS, BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE AND PEPPERS: Apply at first sign of disease infestation. Spray at 7-day intervals following rain, or as needed to control powdery mildew. Can be used up to day of harvest. **STRAWBERRIES:** Apply at first sign of disease infestation. Repeat at 3-week intervals to control powdery mildew. Do not use on strawberry fruit that will be used for canning. Can be used up to the day before harvest. **APPLES & PEARS:** Apply at pre-bloom stage or after petal fall as needed to control powdery mildew. NOTE: Some apple and pear varieties are sulfur sensitive; test for varietal sensitivity prior to full-scale treatment. **CITRUS:** Do not apply within 4 weeks following use of an oil spray. **POTATOES:** Apply before disease infestation appears and repeat as necessary to control powdery mildew.

CUCUMBERS AND SQUASH: Apply at first sign of disease infestation and repeat as necessary to control powdery mildew. Do not spray young plants. **GRAPES:** For control of powdery mildew, apply when new shoots are 6-10 inches. Repeat before blooms open and at 14-day intervals as needed. Note: Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord and other labrusca varieties.

BLACKBERRIES, BLUEBERRIES, RASPBERRIES, LOGANBERRIES AND CURRANTS: Apply each week starting at first bloom color to control powdery mildew. Continue to spray at 10 to 14 day intervals until fruit is set, then repeat as needed or following rain. **MELONS:** Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew. Note: Melon foliage is extremely sensitive to sulfur burn. See paragraph on Use Cautions. **TOMATOES:** Apply at first sign of disease infestation and repeat as necessary to control powdery mildew. Do not spray young plants. **INSECTICIDE AND MITICIDE USE:** Apply at first sighting of insects/mites. Thoroughly moisten top and bottom leaf surfaces where insects/mites are feeding or resting. Apply every 5 to 7 days when insects are present or as needed to prevent damage.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. **Storage:** Store in a cool dry area away from heat or open flame and inaccessible to children. Keep from freezing. **Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Woodstream Corporation and its subsidiaries will not accept liability for injury or damage resulting from product use not in accordance to label instruction. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Distributed By: Safer®, Inc.,
a wholly-owned subsidiary of
Woodstream Corporation

 woodstream

69 North Locust Street,
Lititz, PA 17543
1-800-800-1819 • 717-626-2125
EPA Reg. No. 59913-12
EPA Est. 63416-MN-001

Look for other Safer®
Brand products at
www.saferbrand.com

