

Bon-Neem™ Insecticidal Soap

*Derived from Indian Neem Tree Seeds
and other sources*

R · e · a · d · y · t · o · U · s · e



Aphid



Whitefly



Mite

ACTIVE INGREDIENT	
Potassium Salts of Fatty Acids.....	1.50%*
INERT INGREDIENTS	
Total	98.50%
Total	
EPA Est. No. 4-NY-1	100.00%
*0.32% derived from Neem Seed Oil fatty acids.	
EPA Reg. No. 70191-02-4	

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

(See Back Panel for Additional Precautionary Statements)

TRUSTED SINCE 1926

BONIDE®

- For indoor and outdoor ornamentals and food plants, shrubs, flowers, home garden and greenhouse, fruits, nuts and vegetables
- Specially formulated spray that kills Aphids, Mites, Whiteflies, Adelgids, Caterpillars, Flea Beetles, Scale Insects, Psyllids, Mealy Bugs, Earwigs and many other listed insects.

BON-NEEM INSECTICIDAL SOAP

Ready To Use

Kills: Aphids, Mites, Whiteflies, Adelgids, Caterpillars, Flea beetles, Scale insects, Psyllids, Mealybugs, Earwigs, and many other listed insects.

Use on Listed: Indoor and Outdoor Ornamentals, Food Plants, Shrubs, Flowers, Home Gardens and Green Houses, Fruits, Nuts, and Vegetables.

Store and transport in an upright position.

EPA Est. No. 4-NY-1 EPA Reg. No. 70191-02-4

Buyers Guarantee Limited to Label Claims.

Distributed by
BONIDE PRODUCTS, INC.
6301 Suttiff Road
Oriskany, NY 13424

80

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

This blend of potassium soaps derived from Indian Neem tree seed and other sources is specially formulated as a Ready-to-Use spray that kills mite and insect pests on contact. Target pests include adelgids, aphids, cabbageworms & other caterpillars, earwigs, fleabeetles, lace bugs, leafhoppers, mealy bugs, psyllids, rose slugs, scale insects, sawfly larvae, spider mites, spittle bugs, squashbugs, stinkbugs, tent caterpillars, thrips, and whiteflies on indoor and outdoor ornamental and food plants, shrubs, trees, foliage plants, flowers, home garden and greenhouse fruits, nuts and vegetables.

SHAKE WELL BEFORE USE. Contact and kill listed pests with spray when they first appear or as soon as damage is noticed. Spray infested plant surfaces thoroughly, including lower leaf surfaces, to point of runoff. Re-spray in one week if pests persist. For aphid and mealybug control, re-spray after 2 or 3 days to control high populations. Product can be used up to day of harvest on food plants. Cover aquaria before spraying.

FOR USE ON HOUSEPLANTS, such as ornamental foliage, flowering and food plants in homes, home greenhouses or interior plantscapes.

FOR USE ON FLOWERS, BEDDING PLANTS & PERENNIALS, such as those growing OUTDOORS or around the home.

FOR USE ON WOODY LANDSCAPE SHRUBS, & TREES, including deciduous and evergreen species such as dogwood, crabapple, crape myrtle, lilac, holy, and conifers.

FOR USE ON HOMEGROWN FRUITS & NUTS, such as apples, blackberries, cherries, cranberries, English walnuts, grapes, grapefruit, lemons, melons, nectarines,

Page: One

oranges, peaches, pears, pecans, plum, raspberries, strawberries, and many Exotic, Semitropical or Tropical fruiting plants.

FOR USE ON HOMEGROWN VEGETABLES, such as asparagus, beans, beets, broccoli, cabbage (& other cole crops), celery (and other leafy vegetables), cucumber (& other cucurbits), eggplant, herbs, lettuce, okra, peas, peppers, potatoes, radish, spices (various), spinach, squash & tomatoes.

USE RESTRICTIONS: Do not use on newly transplanted or drought stressed plants or on BLEEDING HEART, CROWN OF THORNS, GARDENIA, HORSE CHESTNUT, JADE PLANT, JAPANESE MAPLE, LANTANA, MOUNTAIN ASH OR SWEETPEAS, or tender new growth of conifers. Product use may discolor foliage of spruce or other conifers. Surface marking may result from presence of sooty molds on foliage or fruits of some plants, particularly citrus and some varieties of pome fruits.

Test a few inconspicuous leaves of BALD CYPRESS, BEGONIAS, CAMELLIAS, CHRYSANTHEMUMS, CUCURBITS, DELICATE FERNS, EUPHORBIA, FUCHSIAS, GRAPE, IMPATIENS, IVIES, PALMS, PEPPERS, POINSETTIAS, NARROW LEAF EVERGREENS, SCHEFFLERA AND SUCCULENTS, and flower of AFRICAN VIOLET, AGERATUM, AZALEA, BEGONIA, CHRYSANTHEMUM, DAHLIA, GERANIUM, IMPATIENS, LILY, MARIGOLD, PANSY, PETUNIA, SALVIA, SNAPDRAGON, VINCA AND ZINNIA for sensitivity before application (or reapplication) to entire plant. If wilting of leaves or flowers occurs within a few hours, rinse test plants with clean water.

PRECAUTIONARY STATEMENTS

Hazards to humans and domestic animals: CAUTION - Causes moderate eye and skin irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Page: Three

Page: Two

STATEMENT OF PRACTICAL TREATMENT: If in eyes - Hold eyelids open and flush with a steady gentle stream of water for 15 minutes. Seek medical attention if irritation persists. **If on skin -** Wash with plenty of soap and water. Seek medical attention if irritation persists.

Environmental Hazards: This product may be hazardous to aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

Bee Hazard: This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

STORAGE AND DISPOSAL

STORAGE: Store in cool place (preferably in a locked storage area) and keep container tightly sealed. Avoid contamination of feed or foodstuffs.

DISPOSAL: PESTICIDE DISPOSAL - Dispose of unused product by securely wrapping original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse before discarding in trash.

Page: Four